

PEERING THROUGH THE STELLAR WIND OF IGR J19140+0951 USING RXTE AND INTEGRAL OBSERVATIONS

Prat, L.¹, Rodriguez, J.¹ and Hannikainen, D.C.²

Abstract. IGR J19140+0951 was discovered by the satellite *INTEGRAL* in 2003. This is a High Mass X-ray binary, in which the compact object is deeply embedded in the wind coming from the supergiant companion star. Due to the high level of photoelectric absorption, this type of source couldn't be detected without hard X-rays observations. The spectra obtained in the 3–80 keV range using *RXTE* and *INTEGRAL* have allowed us to perform a precise spectral analysis of the system along its binary orbit. We were able to confirm the supergiant nature of the companion star and the neutron star nature of the compact object. Using a simple stellar wind model to describe the evolution of the photoelectric absorption, we were able to restrict the orbital inclination angle to the range 37–75°. Finally, we have detected a so-called "soft excess" in at least four observations, for the first time for this source. Such soft excesses have been reported in several HMXBs in the past. We discuss the possible origin of this excess, and suggest, based on its spectral properties and occurrences close to the superior conjunction, that it may be explained as the reprocessing of the X-ray emission originating from the neutron star by the surrounding ionized gas.

1 Introduction

High Mass X-ray Binaries (HMXBs) are binary systems consisting of a compact object orbiting a massive companion star. Prior to the launch of the International Gamma-ray Astrophysics Laboratory (*INTEGRAL*) in 2002, most of the known HMXBs contained a Be companion. In Be-type HMXBs, the compact object emits strong X-ray flashes when it crosses the equatorial plane of the companion star, where a thick disk of matter originating from the stellar wind is present. O and B-type stars have more isotropic stellar winds which absorb the X-ray emission of the compact object, rendering them almost undetectable below a few keV. Thanks to its sensitivity in the soft gamma-ray range, however, *INTEGRAL* has found many such systems in the past few years (see e.g. Liu et al. 2006; Bodaghee et al. 2007). In this perspective, the use of X-ray spectroscopy at different orbital phases makes it possible to probe the stellar wind, providing two-dimensional information on the density and ionization structure of the wind. For instance, the soft excess that is present in the soft X-ray spectra of many HMXBs, whose origin is still quite mysterious, is linked to the physics of the wind close to the compact object, especially the region where the fast moving stellar wind collides with the slow moving and highly ionized gas surrounding the compact object (Hickox et al. 2004).

IGR J19140+0951 was discovered by *INTEGRAL* in March 2003 (Hannikainen et al. 2003). Prior to our study, it was identified as a HMXB with an orbital period of 13.552 days (Wen et al. 2006), likely composed of a neutron star (Rodriguez et al. 2005) orbiting a B0.5 I type supergiant companion star (Hannikainen et al. 2007). Thus, it is a good candidate to study the properties of the obscured X-ray binaries.

2 Spectral analysis

Our data cover the period March 2004–September 2007 with 32 observations taken simultaneously by the two satellites. They were reduced using the latest software packages available (for more details on the analysing process and on the results presented here, we refer to Prat et al. 2008). Since IGR J19140+0951 is a fairly faint

¹ DSM/IRFU/Service d'Astrophysique, CEA/Saclay, F-91191 Gif-sur-Yvette, France

² Metsähovi Radio Observatory, Helsinki University of Technology TKK, Metsähovintie 114, FI-02540 Kylmäla, Finland




Fig. 1. Left : Model absorption (continuous line) and experimental absorption (Δ symbols), for the stellar model with parameters $M_{\star} = 20M_{\odot}$, $R_{\star} = 21R_{\odot}$ and $\beta = 0.5$. Phase 0 corresponds to when the compact object is located between the Earth and the companion star (inferior conjunction). **Right :** Most probable orbital inclination, as a function of the companion radius, for a typical set of parameters. The + symbols correspond to the most probable orbital inclination, confidence contours are drawn at the 25, 68 and 90% confidence levels. The dotted line is the “eclipse limit”: points above this line are excluded as it would imply an eclipse of the neutron star by its companion, which is not observed.

source, we also corrected the spectra for the Galactic X-ray background (GXB), using the parameters measured by Valinia & Marshall (1998). This correction aimed mostly at minimizing the errors on the measure of the photoelectric absorption, by removing the Galactic contribution and keeping only the absorption intrinsic to the source.

We tested several models in analysing the spectra, starting with phenomenological models. The data were well fitted using an absorbed power law combined with a high energy cut-off. An iron fluorescent line at 6.4 keV is present in the observations where the source is detected at a high enough significance. Note that the high energy cut-off is only detected in the observations where the source flux is sufficiently high, thus giving enough precision at the high energy end of the spectra. The presence of this cut-off allowed us to identify the emission process as thermal Comptonization, which is quite common for this type of source. This is also confirmed by the good fitting results obtained with the more physical Comptonization model `COMPTT`, in XSPEC notations.

The photoelectric absorption N_{H} undergoes strong variations, well correlated with the orbital phase (Fig. 1, left). Since the orbit of the X-ray source is probably almost circular, one possible explanation for the observed change in N_{H} relies on an inclined orbit for the system. Qualitatively, when the neutron star is behind its companion its light has to travel a longer distance through a denser part of the stellar wind, so the absorption in the soft X-rays increases. Therefore, at the superior conjunction we should observe a maximum in the absorption, which is indeed the case.

The unabsorbed flux undergoes strong variations that are not correlated with the orbital phase; this may be a consequence of the details of the accretion process. The stellar wind is not perfectly homogeneous and the accretion onto the neutron star, following the magnetic field line to the magnetic poles, is a highly unstable and complicated process (see e.g. Takata et al. 2006, and references therein, for a review of the emission processes). Since our observations are spread over several years, it is not surprising that the source flux is highly variable. This may also be a direct consequence of the clumpiness of the stellar wind: depending on the times of the observation, the compact object may be in the course of accreting a denser portion of the wind (Blay et al. 2008). Unfortunately, the rather short exposure times and the too long periods between successive observations restrained precise characterisations of this clumpiness.

3 Soft Excess detection

But a more interesting characteristic is the detection of a so called “soft excess”. During our study, some spectra exhibited an excess in the soft X-ray part of the spectra, which we modeled by adding a black body component to the model. This feature has already been observed in many X-ray binaries (see Hickox et al. 2004, for a review). Its origin is mysterious, but can be linked to the immediate environment of the neutron star, and explained qualitatively by the following scenario. The stellar wind of massive stars is mostly accelerated by bound-bound transitions of atoms. The neutron star, due to its high energy emission, ionizes the surrounding material, so the already ionized gas around the neutron star is no longer accelerated by the stellar radiation field. When the compact object moves along its orbit, the hot gas will gradually be overtaken by the stellar wind. This will lead to the formation of a “tail” trailing the neutron star.

As the gas in this tail is ionized, its absorption will be lower, but its precise effect on the emission is not clear. Since the soft excess is only visible around the superior conjunction (between phases 0.40 and 0.75), one possibility is that the hard X-ray emission may be scattered around the tail. The precise study of this effect, however, would require better spectral resolution in the soft X-ray range than what is available on *RXTE*.

4 Wind model

In order to constrain several parameters for the system, we used a simple “ β -law” wind model: it adequately describes the strong stellar winds of B-type stars, which is usually taken to be stationary and spherically symmetric. See e.g. Levine et al. (2004) for a precise description of the method. Using the evolution of the photoelectric absorption, we were able to constrain the inclination of the system and the mass-loss rate of the companion star. For the most probable stellar radius of $21 R_{\odot}$, the lower inclination limit is constrained between 37 and 42° (Fig. 1, right), with χ^2 being at a minimum in the range 55 – 63° . Fig. 1, left, shows the best-fit model against the experimental normalized data, with a good agreement. The mass-loss rate, \dot{M}_{\star} , is constrained to the range 0.6 – $1.0 \times 10^{-7} M_{\odot}/\text{year}$, which is consistent with what is expected for a B0.5 type supergiant.




Fig. 2. Diagram of IGR J19140+0951 as it could be seen from the Earth. The orbital inclination of the system is taken to be $\sim 65^{\circ}$.

Fig. 2 summarizes schematically our main results. We found that photoelectric absorption highly correlates with the orbital phase of the system. Using a simple stellar wind model we found a rather high orbital inclination, $\sim 65^{\circ}$. We have detected a soft excess in some observations, just before the superior conjunction, in the area

indicated on the figure. This may be explained by a cloud of highly ionised gas surrounding the neutron star. Because of its ionisation, this gas is less accelerated by the stellar wind and, while being gradually overtaken by the wind, tends to form a “tail”. This tail could scatter the hard X-ray emission from the compact object and thus explain the soft excess feature. However, these results remain qualitative, as a precise determination of the geometry of the stellar wind would require a better sensitivity in the soft X-ray range, which could be achieved for instance with the use of the *XMM* or *Chandra* satellites.

5 Conclusions

The study of X-ray binaries is challenging since it is often difficult, if not impossible, to identify their visible and infrared counterparts. Even if an infrared counterpart were observed, the distance to some systems prohibits the measurement of the orbital characteristics. Our study shows that X-ray observations can overcome these limitations and produce very precise inferences. The *RXTE* and *INTEGRAL* observations of IGR J19140+0951 have led to good measurements of the orbital period of the system, constraints on its inclination angle, and the diagnostic of the type of the companion star (supergiant O or B)

Moreover, we can use the compact object to probe the stellar wind of the companion. In the case of IGR J19140+0951, we diagnosed the wind density, the probable presence of clumps, and the wind structure around the neutron star. More precise observations could lead to constraints on the mass and radius of the companion, and better constraints on the stellar wind. This allows the study of new *INTEGRAL* sources, either distant or highly absorbed, and ultimately the determination of new useful data for X-ray binary evolution scenarios.

References

- Blay, P., Martnez-Nez, S., Negueruela, I., Pottschmidt, K., Smith, D. M., et al. 2008, astro-ph/08064097
Bodaghee, A., Courvoisier, T. J.-L., Rodriguez, J., et al. 2007, A&A, 457, 585
Hannikainen, D., Rawlings, M. G., Muhli, P., et al. 2007, MNRAS, 380, 665
Hannikainen, D., Rodriguez, J., & Pottschmidt, K. 2003, IAUC, 8088
Hickox, R. C., Narayan, R., & Kallman, T. R. 2004, ApJ, 614, 881
Liu, Q. Z., van Paradijs, J., van den Heuvel, E. P. J. 2006, A&A, 455, 1165
Levine, A. M., Rappaport, S., Remillard, R., & Savcheva, A. 2004, ApJ, 617, 134
Prat, L., Rodriguez, J., & Hannikainen, D. 2008, MNRAS, 389, 301
Rodriguez, J., Cabanac, C., Hannikainen, D., et al. 2005, A&A, 432, L17
Takata J., Shibata, S., Hirotoni, K., & Chang, H.-K. 2006, MNRAS, 366, 1310
Valinia, A., & Marshall, F. E. 1998, ApJ, 505, 134
Wen L., Levine, A., Corbet, R., & Bradt, H. 2006, ApJS, 163, 372